

DISTILLABILITY CRITERION FOR ALL BIPARTITE GAUSSIAN STATES

G GIEDKE*, L-M DUAN, I CIRAC, AND P ZOLLER,

Institut für Theoretische Physik, Universität Innsbruck, Technikerstrasse 25, 6020 Innsbruck, Austria

We prove that all inseparable Gaussian states of two modes can be distilled into maximally entangled pure states by local operations. Using this result we show that a bipartite Gaussian state of arbitrarily many modes can be distilled if and only if its partial transpose is not positive.

Keywords: entanglement, distillation, Gaussian states

Communicated by: S Braunstein and C Fuchs

The existence of *pure* entangled states of two or more systems entails the possibility of finding new applications of Quantum Mechanics, in particular in the fields of computation and communication [1]. In practice, however, systems are exposed to interactions with the environment, that transform pure into *mixed* states, which may no longer be useful for quantum communication. Fortunately, there exist methods to recover pure entangled states from mixed ones in certain situations. These processes are called *entanglement distillation* (or purification) [2], and consist of local operations and classical communication transforming several copies of a mixed entangled state into (approximately) pure entangled states which can then be used for quantum communication. In fact, applying this method in the appropriate way one can construct quantum repeaters [3] that should allow efficient quantum communication over arbitrarily long distances even via a noisy channel.

For this reason it is important to determine whether a given state is distillable or not. In general, the answer to this question is not known. At the moment we only have conditions that are necessary or sufficient for distillability, but not both. Clearly, only inseparable states can be distilled. Moreover, as shown by Horodecki *et al.* [4], there exists a stronger necessary condition, namely that ρ must have non-positive partial transpose (npt). In fact, there are entangled states which are not distillable since their density matrices remain positive under partial transposition [5]. Furthermore, there is evidence that this condition is not sufficient, since there exist npt states that nevertheless seem to be undistillable [6]. The existence of undistillable npt states would have interesting consequences such as non-additivity and non-convexity of the entanglement of formation [7].

On the other hand, a useful sufficient criterion, the so-called reduction criterion [8], has been established. It states that, given a state ρ on the composite Hilbert space $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$, if there exists a vector $|\psi\rangle \in \mathcal{H}$ such that

$$\langle\psi|\operatorname{tr}_B\rho\otimes\mathbb{1}-\rho|\psi\rangle<0. \quad (1)$$

then the state ρ is distillable. Here, tr_B stands for the partial trace with respect to the second

*email: geza.giedke@uibk.ac.at

subsystem. An important aspect of this criterion is that if one can find a state $|\psi\rangle$ satisfying (1), then one can explicitly construct a protocol to distill ρ .

Up until now, nearly all work on the distillability problem has considered states of finite dimensional systems, see [9] for a current overview. In particular it was shown that states systems consisting of one qubit and an N -level, $N \geq 2$ system are distillable if and only if (iff) they are npt [10, 6]. An alternative setting for quantum information processing, which considers infinite dimensional systems [continuous variables (CV) or “modes”] in Gaussian states is receiving increasing attention recently [11, 12]. For CV systems some distillation protocols for particular states have been proposed [13], and the existence of bound entangled states has been proved [14, 15], but the question of distillability in general has not been addressed.

In this article we answer this question completely for all Gaussian states. We will prove that

Theorem 1 (*Distillability Criterion*)

A Gaussian state of $N \times M$ modes is distillable if and only if its partial transpose is negative.

This shows that there are no npt bound entangled Gaussian states and, in particular, that for systems of $1 \times N$ modes all entangled states (npt is necessary for inseparability of such systems [15, 16, 17]) are distillable and thus useful for quantum communication. Moreover, our proof, which is based in part on the reduction criterion, provides an explicit protocol that accomplishes distillation for all those states. After introducing the necessary notation and properties of Gaussian states, the remainder of the paper is devoted to the proof of Theorem 1.

We consider bipartite systems composed of two subsystems, A and B, which consist of N and M “modes” [distinguishable infinite dimensional quantum systems with Hilbert space $L^2(\mathbb{R})$], respectively. The joint system is referred to as a “ $N \times M$ system”. It is convenient to describe the state ρ of such a system by its characteristic function (e.g., [18])

$$\chi(x) = \text{tr}[\rho D(x)]. \quad (2)$$

Here $x = (q_1, p_1, \dots, q_{N+M}, p_{N+M}) \in \mathbb{R}^{2N+2M}$ is a real vector and

$$D(x) = e^{-i \sum_k (q_k X_k + p_k P_k)}, \quad (3)$$

where X_k and P_k are operators satisfying the canonical commutation relations ($\hbar = 1$). A characteristic function χ uniquely defines a state ρ_χ . In the following we exclusively consider *Gaussian states*, i.e. states for which χ is a Gaussian function of x [19]

$$\chi(x) = e^{-\frac{1}{4}x^T \gamma x - id^T x}, \quad (4)$$

where γ is the *correlation matrix* (CM) and $d \in \mathbb{R}^{2N+2M}$ the *displacement*. Thus, a Gaussian state is fully characterized by its CM γ and displacement d . These states are of particular interest, since they comprise essentially all CV states that can be prepared in the lab with current technology.

A matrix γ is the CM of a physical state iff (e.g. [22]) it is strictly positive, real, symmetric $2(N+M) \times 2(N+M)$ and satisfies

$$\gamma \geq J^T \gamma^{-1} J, \quad (5)$$

where $J_{N+M} = \bigoplus_{k=1}^{N+M} J_1$ [20] with $J_1 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

Simon [17] noted that for CV states partial transposition is equivalent to the orthogonal transformation $\Lambda_B(q_A, p_A, q_B, p_B) = (q_A, p_A, q_B, -p_B)$ on phase space, i.e., the momentum coordinates referring to B are inverted. For a Gaussian state this means that its CM is changed to $\tilde{\gamma} = \Lambda_B \gamma \Lambda_B$ and the displacement to $\Lambda_B d$. A Gaussian state with CM γ has negative partial transpose (npt) iff $\tilde{\gamma}$ does not satisfy Ineq. (5) [17, 15], or, equivalently, iff

$$\gamma \not\geq \tilde{J}^T \gamma^{-1} \tilde{J}, \quad (6)$$

where $\tilde{J} = \Lambda_B J \Lambda_B^T$ is the “partially transposed” J in which the J_1 ’s corresponding to B’s modes are replaced by $-J_1$.

The first part of the proof of the theorem is concerned with the special case of a bipartite two-mode Gaussian state: $N = M = 1$. Any such state can be transformed into what we called the *standard form*, using local unitary operations only [16, 17]. For a state in standard form the displacement $d = 0$ and the CM γ has the simple form

$$\gamma = \begin{pmatrix} A & C \\ C^T & B \end{pmatrix}, \quad (7)$$

where

$$A = \begin{pmatrix} n_a & 0 \\ 0 & n_a \end{pmatrix}, B = \begin{pmatrix} n_b & 0 \\ 0 & n_b \end{pmatrix}, C = \begin{pmatrix} k_x & 0 \\ 0 & k_p \end{pmatrix}. \quad (8)$$

The local unitaries needed to achieve this form are linear Bogoliubov transformations, i.e., generated by Hamiltonians that are at most quadratic in the operators $X_{1,2}, P_{1,2}$. The four real parameters (n_a, n_b, k_x, k_p) fully characterize a 1×1 Gaussian state up to local linear Bogoliubov transformations (LLBT). They can be easily calculated from the four LLBT-invariant determinants $\det A, \det B, \det C$, and $\det \gamma$ as follows:

$$n_a = \sqrt{\det A}, n_b = \sqrt{\det B}, k_x k_p = \det C, \quad (9a)$$

$$(n_a n_b - k_x^2)(n_a n_b - k_p^2) = \det \gamma. \quad (9b)$$

Without loss of generality we choose $k_x \geq |k_p|$. We call a state *symmetric*, if $n_a = n_b = n$, or, equivalently, if $\det A = \det B$.

Now we are prepared for the proof of Theorem 1. We state the three main steps of the proof in three lemmas, which we prove in the remainder of this article.

Lemma 1 (*Distillability of Symmetric 1×1 States*)

A symmetric 1×1 Gaussian state with non-positive partial transpose is distillable.

Lemma 2 (*Symmetrization of 1×1 States*)

Every 1×1 Gaussian state with non-positive partial transpose can be locally transformed into a symmetric npt state.

Lemma 3 (*Concentrating Inseparability in two Modes*)

Every $N \times M$ Gaussian state with non-positive partial transpose can be locally transformed into a 1×1 npt state.

Proof of Theorem 1: The “only if”-part of the Theorem was proven proven by the Horodeckis in [4]. The “if”-part is clearly implied by these three Lemmas, since by Lemma 3 the $N \times M$ case can be reduced to the 1×1 case, and that case by Lemma 2 to the symmetric case. ■

For the proof of Lemmas 1 and 2, it is useful to re-express the conditions (5,6) for 1×1 states in terms of the parameters (9). We find that γ is CM of a physical state iff

$$(n_a n_b - k_x^2)(n_a n_b - k_p^2) + 1 \geq n_a^2 + n_b^2 + 2k_x k_p, \quad (10a)$$

$$n_a n_b - k_x^2 \geq 1, \quad (10b)$$

and that γ is CM of an inseparable (or, equivalently, npt) state, iff in addition it holds that

$$(n_a n_b - k_x^2)(n_a n_b - k_p^2) + 1 < n_a^2 + n_b^2 - 2k_x k_p. \quad (11)$$

Proof of Lemma 1: For this we use that a state is distillable, if there exists a pure state $|\psi\rangle$ such that Ineq. (1) holds. This condition was proved in [8] to be sufficient for distillability of finite dimensional systems. Its extension to infinite dimensions is straightforward: and proved in the appendix.

We show now that for any symmetric npt Gaussian state ρ Ineq. (1) is satisfied with $|\psi\rangle$ taken as the pure two-mode squeezed state $|\psi\rangle = \frac{1}{\cosh r} \sum_n \tanh^n r |nn\rangle$ for sufficiently large $r > 0$. Note that $|\psi\rangle$ is a symmetric Gaussian state in standard form. We denote its CM by γ_ψ and the four parameters (9) are $n_a = n_b = \cosh 2r$, $k_x = -k_p = \sinh 2r$. Let γ_ρ denote the CM of ρ . With these choices, Ineq. (1) becomes [21]

$$2 [\det(\gamma_{\text{tr}_B \rho} + \gamma_{\text{tr}_B \psi})]^{-1/2} - 4 [\det(\gamma_\rho + \gamma_\psi)]^{-1/2} < 0. \quad (12)$$

In the limit of large r (keeping only the leading terms in e^r) this becomes after some simple algebra

$$(n - k_x)(n + k_p) < 1. \quad (13)$$

But Ineq. (13) is implied by the inseparability criterion for symmetric states: if $n_a = n_b = n$ then Ineq. (11) simplifies to

$$|n^2 - k_x k_p - 1| < n(k_x - k_p). \quad (14)$$

For inseparable states we observe [17] that $k_x k_p < 0$, which together with Ineq. (10b) implies that the LHS of Ineq. (14) is equal to $n^2 - k_x k_p - 1$ which can be transformed to $(n - k_x)(n + k_p) + n(k_x - k_p) - 1$ from which Ineq. (13) follows immediately. ■

Since the local operation that will be shown to achieve symmetrization involves a measurement, it is more convenient to describe the state here by its *Wigner function* [18]. It is related to the characteristic function by symplectic Fourier transformation and thus is Gaussian for Gaussian states. The Wigner CM γ_W is related to the (characteristic) CM by $\gamma_W = J^T \gamma^{-1} J$. We denote the four LLBT-invariant parameters for the Wigner CM [which are defined as in (9)] by (N_a, N_b, K_x, K_p) . We use the following easily checked facts: just as the standard form of γ , the standard form of γ_W can be obtained by LLBTs. A state is symmetric iff $N_a = N_b$. The conditions (10,11) can be formulated equivalently in terms of the parameters

(N_a, N_b, K_x, K_p) . While (10a, 11) are identical for the Wigner parameters, in (10b) “ $\geq$ ” is changed to “ $\leq$ ”. We refer to these conditions for the Wigner parameters as (10W, 11W) in the following.

Proof of Lemma 2: If the state is not symmetric, it means that the reduced state at one of the two sides has larger entropy than the other. This suggests to let a pure state interact with the “hotter” side to cool it down. This must be done without destroying the entanglement of ρ . We proceed as follows: ρ is transformed to its Wigner standard form with parameters (N_a, N_b, K_x, K_p) . Now assume that $N_b < N_a$, i.e., B is the hotter side [24]. Take an ancilla mode in the vacuum state and couple it to B’s mode by a beam splitter [23] with transmittivity $\cos^2 \theta$. After a measuring the ancilla’s X -operator [25] results a state $\tilde{\rho}$ with Wigner CM $\tilde{\gamma}_W$ of the form (7) with

$$\tilde{A} = \frac{1}{\nu} \begin{pmatrix} c^2 N_a + s^2 D_x & 0 \\ 0 & c^2 N_a + s^2 N_a N_b \end{pmatrix},$$

$$\tilde{B} = \frac{1}{\nu} \begin{pmatrix} N_b & 0 \\ 0 & [c^2 N_b + s^2] \nu \end{pmatrix}, \quad \tilde{C} = \frac{1}{\nu} \begin{pmatrix} c K_x & 0 \\ 0 & c K_p \nu \end{pmatrix},$$

where the abbreviations $c = \cos \theta, s = \sin \theta, \nu = s^2 N_b + c^2$, and $D_{x,p} = N_a N_b - K_{x,p}^2$ were used. The condition for symmetry, $|\det \tilde{A}| = |\det \tilde{B}|$, requires

$$\tan^2 \theta = \frac{N_a^2 - N_b^2}{N_b - D_x N_a}. \quad (15)$$

Checking (11W) for $\tilde{\gamma}_W$ one sees that the inequality is just multiplied by $(N_b \tan^2 \theta + 1)^{-1} > 0$; therefore the transformed state is inseparable iff the original one was inseparable. It remains to show that there always exists a θ to satisfy (15), i.e., that the right hand side of Eq. (15) is positive. The numerator is positive since we have chosen $N_b < N_a$, the denominator is positive since $N_b < N_a$ and the second part of condition (10W) imply that $(N_a - D_x N_b) > 0$ and the first part of (10W) assures that $(N_a - D_x N_b)(N_b - D_p N_a) \geq (N_a K_x + N_b K_p)^2 \geq 0$, hence all Gaussian states in Wigner standard form can be symmetrized this way. But since every Gaussian state can be brought into Wigner standard form by local unitaries, this completes the proof of Lemma 2. ■

To finish the proofs, we now turn to the general case of $N \times M$ modes. Let γ be the CM of a npt state.

Proof of Lemma 3: The condition (6) is equivalent to $\gamma \not\geq i\tilde{J}$ [15]. Hence, for every npt state with CM γ there exists a vector $z \in \mathbb{C}^{2(N+M)}$ such that for some $\epsilon > 0$

$$z^\dagger (\gamma - i\tilde{J}) z \leq -\epsilon < 0. \quad (16)$$

The idea of the proof is that γ can be locally transformed such that at both sides all but one mode can be discarded, and the resulting (reduced) 1×1 state is still npt. Then it is distillable by Lemmas 1 and 2.

Write z in Eq. (16) as $z = z^{(A)} \oplus z^{(B)}$ with real and imaginary parts $z_r^{(x)}, z_i^{(x)}$, ($x = A, B$). We can always find a z such that $z_r^{(x)}$ and $z_i^{(x)}$ are not skew-orthogonal, i.e. $(z_r^{(x)})^T J z_i^{(x)} \neq 0$ for both $x = A, B$ [26].

Now we have to find two symplectic transformations $S_x, x = A, B$ that map the span of $\{z_r^{(x)}, z_i^{(x)}\}$ to the span of $\{e_1, e_2\}$, where $e_1 = (1, 0, \dots, 0)$ etc. After performing the local transformation $S = S_A \oplus S_B$, both A and B can discard all but the first mode of their systems and still have an npt entangled (and thus distillable) state. That such symplectic transformations S_A, S_B always exist is seen as follows: let $f_1 = z_r, f_2 = -z_i/(z_r^T J z_i)$; we can always extend f_1, f_2 into a *symplectic basis* [27] $f_k : k = 1, \dots, 2n$ such that $f_{2k}^T J f_{2l+1} = \delta_{kl}, f_{2k}^T J f_{2l} = 0 = f_{2k+1}^T J f_{2l+1}$. Then S defined by $S e_k = f_k$ is symplectic [27] and S^{-1} maps $\text{span}\{z_r, z_i\}$ to $\text{span}\{e_1, e_2\}$, i.e.

$$\hat{z}^{(x)} := S_x^{-1} z^{(x)} = a_x e_1 + b_x e_2, \quad (17)$$

$a_x, b_x \in \mathbb{C}$. Consequently we have for $\hat{\gamma} := (S_A \oplus S_B)^T \gamma (S_A \oplus S_B)$

$$(\hat{z}^{(A)} \oplus \hat{z}^{(B)})^\dagger (\hat{\gamma} - i\tilde{J})(\hat{z}^{(A)} \oplus \hat{z}^{(B)}) < 0. \quad (18)$$

Using Eq. (17) we see that only the matrix elements $(\hat{\gamma})_{kl}$ with $k, l = 1, 2, N+1, N+2$ contribute to the lhs of Eq. (18). Thus Ineq. (18) does not change if we replace $\hat{\gamma}$ by the two-mode CM $\hat{\gamma}_{red}$ obtained from $\hat{\gamma}$ by discarding all rows and columns referring to modes other than 1 and $N+1$. This is the CM of the state in which A and B discard all but their first mode each. Thus Ineq. (18) shows that the state ρ_{red} corresponding to $\hat{\gamma}_{red}$ is npt. But ρ_{red} is a two-mode state and thus distillable by the first part of the proof. ■

Note that all the operations needed to transform a general $N \times M$ npt state into a symmetric 1×1 entangled state can be implemented quantum optically with current technology: they require nothing but squeezers, beam splitters, phase shifters [23], homodyne measurements, and the discarding of subsystems. Once a state has been transformed to symmetric standard form, the protocol of Ref. [8] can be used to obtain maximally entangled states in a finite dimensional Hilbert space. While a *practical* distillation protocol for such Gaussian states remains to be found (see however [13]), it is worth noting, that the main part of the universal protocol of [8], namely the filtering operation and the joint measurement, are for symmetric Gaussian states implemented by the procedure of Duan *et al.*[13]; for details see [12, ch. II.8].

In conclusion, we have answered the distillability question for all Gaussian states: such states are distillable if and only if they are npt. In particular, all entangled Gaussian states of $1 \times N$ modes are distillable, and there exist no npt bound entangled Gaussian states.

Acknowledgements

G.G. acknowledges financial support by the Friedrich-Naumann-Stiftung. This work was supported by the Austrian Science Foundation (SFB “Control and Measurement of Coherent Quantum Systems”, Project 11), the EU (TMR network ERB-FMRX-CT96-0087 and the project EQUIP, contract IST-1999-11053), the ESF, and the Institute for Quantum Information GmbH, Innsbruck.

References

1. C.H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W.K. Wootters, *Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels*, Phys. Rev. Lett., **70**, 1895 (1993);

- A. Ekert, *Quantum cryptography based on Bell's theorem*, Phys. Rev. Lett. **67**, 661 (1991).
2. C.H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J.A. Smolin, and W.K. Wootters, *Purification of Noisy Entanglement and Faithful Teleportation via Noisy Channels*, Phys. Rev. Lett. **76**, 722 (1996); quant-ph/9511027.
- C.H. Bennett, D.P. DiVincenzo, J.A. Smolin, and W.K. Wootters, *Mixed-state entanglement and quantum error correction*, Phys. Rev. A **54** 3824 (1996); quant-ph/9604024.
3. H.-J. Briegel, W. Dür, J.I. Cirac, and P. Zoller, *Quantum repeaters: The role of imperfect local operations in quantum communication*, Phys. Rev. Lett. **81**, 5932 (1998); quant-ph/9803056.
4. M. Horodecki, P. Horodecki, and R. Horodecki, *Mixed-state entanglement and distillation: Is there a "bound" entanglement in nature?*, Phys. Rev. Lett. **80**, 5239 (1998).
5. P. Horodecki, *Separability Criterion and inseparable mixed states with positive partial transpose*, Phys. Lett. A **232**, 333 (1997).
6. W. Dür, J.I. Cirac, M. Lewenstein, D. Bruß, *Distillability and partial transposition in bipartite systems*, Phys. Rev. A **61**, 062313 (2000), quant-ph/9910022.
- D.P. DiVincenzo, P.W. Shor, J.A. Smolin, B.M. Terhal, A.V. Thapliyal, *Evidence for bound entangled states with negative partial transpose*, Phys. Rev. A **61**, 062312 (2000); quant-ph/9910026.
7. P.W. Shor, J.A. Smolin, B.M. Terhal, *Nonadditivity of Bipartite Distillable Entanglement follows from Conjecture on Bound Entangled Werner States*, Phys. Rev. Lett. **86**, 2681 (2001); quant-ph/0010054.
8. M. Horodecki, P. Horodecki, *Reduction criterion of separability and limits for a class of distillation protocols*, Phys. Rev. A **59**, 4206 (1999).
9. M. Lewenstein, D. Bruß, J.I. Cirac, B. Kraus, M. Kuś, J. Samsonowicz, A. Sanpera, and R. Tarrach, *Separability and distillability in composite quantum systems – a primer*, J. Mod. Opt. **47**, 2481 (2000), quant-ph/006064.
10. M. Horodecki, P. Horodecki, and R. Horodecki, *Inseparable Two Spin-1/2 Density Matrices Can Be Distilled to a Singlet Form*, Phys. Rev. Lett. **78**, 574 (1997).
11. L. Vaidman, *Teleportation of quantum states*, Phys. Rev. A **49**, 1473 (1994).
- S.L. Braunstein and H.J. Kimble, *Teleportation of Continuous Quantum Variables*, Phys. Rev. Lett. **80**, 869 (1998);
- A. Furusawa, J.L. Sørensen, S.L. Braunstein, C.A. Fuchs, H.J. Kimble, and E.S. Polzik, *Unconditional Quantum Teleportation*, Science **282**, 706 (1998).
12. S.L. Braunstein and A.K. Pati (eds.), *Quantum Information Theory with Continuous Variables*, Kluwer Academic Publishers, Dordrecht 2001, in press.
13. T. Opatrny, G. Kurizki, and D.-G. Welsch, *Improvement on teleportation of continuous variables by photon subtraction via conditional measurement*, Phys. Rev. A **61**, 032302 (1999); quant-ph/9907048;
- S. Parker, S. Bose, and M.B. Plenio, *Entanglement quantification and purification in continuous variable systems*, Phys. Rev. A **61**, 032305 (1999); quant-ph/9906098;
- L.-M. Duan, G. Giedke, J.I. Cirac, and P. Zoller, *Entanglement Purification of Gaussian Continuous Variable Quantum States*, Phys. Rev. Lett. **84**, 4002 (2000); quant-ph/9912017.
14. P. Horodecki, M. Lewenstein, *Bound Entanglement and Continuous Variables*, Phys. Rev. Lett. **85**, 2657 (2000); quant-ph/0001035.
15. R.F. Werner and M.M. Wolf, *Bound entangled Gaussian States*, Phys. Rev. Lett. **86**, 3658 (2001); quant-ph/0009118.
16. L.-M. Duan, G. Giedke, J.I. Cirac, and P. Zoller, *Inseparability criterion for continuous variable systems*, Phys. Rev. Lett. **84**, 2722 (2000); quant-ph/9908056.
17. R. Simon, *Peres-Horodecki separability criterion for continuous variable systems*, Phys. Rev. Lett. **84**, 2726 (2000); quant-ph/9909044.
18. C.W. Gardiner and P. Zoller, *Quantum Noise* 2nd ed., Springer-Verlag, Berlin (1999).
19. J. Manuceau and A. Verbeure, *Comm. Math. Phys.* **9**, 293 (1968).
20. For convenience we use direct sum notation for matrices and vectors. That is, if A and B are

- $n \times n$ and $m \times m$ matrices, resp., then $A \oplus B \in M_{n+m, n+m}$ is a block diagonal matrix of blocks A and B . Similarly, if $f_1 \in \mathbb{R}^n$ and $f_2 \in \mathbb{R}^m$ are two vectors, then $f_1 \oplus f_2 \in \mathbb{R}^{n+m}$ is a vector whose first n components are given by the entries of f_1 and the last m by those of f_2 .
21. H. Scutaru, *Transition Probabilities for Quasifree States*, J. Math. Phys. **39**, 6403 (1998).
 22. H. Scutaru, *The states with Gaussian Wigner function are quasi-free states*, Phys. Lett. A **141**, 223 (1989).
 23. We use quantum optical terminology for the following LBT: a beam splitter of transmittivity t coupling the modes k and l refers to the unitary generated by $t(P_k X_l - X_l P_k)$, a phase shifter for the k th mode is generated by $X_k^2 + P_k^2$, a squeezing transformation by $X_k P_k + P_k X_k$, and a displacement by $aX_k + bP_k$.
 24. Note that since γ_W is essentially the inverse of γ , large (Wigner parameter) N_a implies *small* entropy of the reduced state, while large (characteristic function parameter) n_a implies large entropy of the reduced state; more precisely we have: $N_a = n_b / \sqrt{\det \gamma}$, $N_b = n_a / \sqrt{\det \gamma}$.
 25. If a measurement of an observable X_k or P_k is performed, the Wigner function of the final state of the remaining modes is calculated as follows: replace the phase space coordinate referring to the measured operator by the measured result and integrate out the conjugate phase space coordinate. Note that the CM of final state is independent of the measurement outcome.
 26. In case the original vectors z_r, z_i are skew-orthogonal, we can replace z_i by $z'_i = z_i + \delta J z_r$ where δ is so small that Ineq. (16) still holds for $z' = z_r + i z'_i$.
 27. V.I. Arnold, *Mathematical Methods of Classical Mechanics* 2nd. ed., Springer Verlag, New York, 1989.

Appendix A

We show that Ineq. (1) implies distillability even for $\dim \mathcal{H} = \infty$. Let $\{|k\rangle : k = 0, 1, \dots\}$ be an orthonormal basis of $\mathcal{H}$, let $\mathcal{H}_n = \text{span}\{|0\rangle, |1\rangle, \dots, |n\rangle\}$, $P_{\mathcal{H}_n}$ the orthogonal projector on $\mathcal{H}_n$, and let ρ be a density matrix on $\mathcal{H} \otimes \mathcal{H}$. Let $\mathcal{E}(\rho) = \text{tr}_B \rho \otimes \mathbb{1} - \rho$ be the map occuring on the lefthand side of (1). Assume that $\exists |\psi\rangle \in \mathcal{H}, \epsilon > 0$ such that $\langle \psi | \mathcal{E}(\rho) | \psi \rangle < -\epsilon < 0$. Since $\rho_n = P_{\mathcal{H}_n} \rho P_{\mathcal{H}_n}$ converges in the weak topology to ρ , there is $N \geq 0$ such that $\langle \psi | \mathcal{E}(\rho_n) | \psi \rangle < -\epsilon/2$ for all $n \geq N$. Thus ρ can be projected by local operations onto a distillable state ρ_N and is therefore itself distillable. ■